1. Chronic bowel disorders

1.1 Coeliac Disease	Primary Care	Acute/ Specialist Services
Refer to NHS Forth Valley Gluten Free Food List		

1.2 Diverticular disease and diverticulitis	Primary Care	Acute/ Specialist Services
No specific formulary choices		

1.3 Inflami	matory bowel disease	Primary Care	Acute/ Specialist Services
Aminosalic	ylates		
	Balsalazide Sodium	 	✓
	Mesalazine (Octasa [®] , Salofalk [®] , Pentasa [®])	\$	✓
	Oral mesalazine products should be prescribed by brand name only due between different brands.	e to differences	in
	Sulfasalazine	\$	✓
Corticoster	oids		
	Budesonide (Budenofalk [®] , Cortiment [®] , Entocort [®] , Jorveza [®])	+	✓
Jorveza® oro-	Note different brands of budesonide have different licensed indications. dispersible tablets are approved for use for the treatment and maintena besophagitis where a minimum 4 week trial of a PPI has been ineffective		
	Prednisolone (foam enema, rectal solution, suppositories)	+	✓
Immunosu	pressants		
	Adalimumab (Imraldi [®])		✓
	Azathioprine	 	✓
	Ciclosporin	\$	✓
	Infliximab (Remsima®)		✓
	Mercaptopurine	\$	✓
	Methotrexate	+	✓
	Tofacitinib		✓

Key \checkmark Initiate and continue \div Continue where appropriate.

	Vedolizumab (infusion, s.c.)		✓
Comme	t: Biologic therapies can only be prescribed by a consultant gastroenterolo	gist.	

1.4 Irrita	ble Bowel Syndrome	Primary Care	Acute/ Specialist Services	
Antispasi	modics			
	Peppermint Oil	✓	✓	
Guanylate	Guanylate Cyclase-C Receptor Agonists			
	Linaclotide (Constella®)	\$	✓	
Comment: To be prescribed in line with <u>SMC advice</u>				

1.5 Short bowel syndrome	Primary Care	Acute/ Specialist Services
No Formulary Choices		

2 Constipation and bowel cleansing	Primary Care	Acute/ Specialist Services
Bowel cleansing solutions		
Moviprep [®]		✓
Picolax [®]		✓

2.2 Con	stipation	Primary Care	Acute/ Specialist Services
Bulk-forn	ning laxatives		
	Ispaghula Husk	✓	✓
Osmotic	Laxatives		
	Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride (Laxido [®] , Laxido Paediatric [®])	✓	✓
	Lactulose	✓	✓
	: Lactulose may take up to 48 hours to act and is therefore unsuitable to prescribing.	for relief of acute	symptoms

Key \checkmark Initiate and continue \Leftrightarrow Continue where appropriate.

Fleet [®] Ready-to-use Enema	✓	✓		
Phosphate enema	✓	✓		
Softening laxatives				
Docusate	✓	✓		
Stimulant laxatives				
Senna (1 st line)	✓	✓		
Bisacodyl (2 nd line)	✓	✓		
Co-danthramer (palliative care use only)	✓	✓		
Glycerol suppositories	✓	✓		
Comment: Prolonged use of stimulant laxatives is not recommended.				
Opioid receptor antagonists				
Naloxegol (Moventig [®])	\$	✓		

3. Diarrh	oea	Primary Care	Acute/ Specialist Services	
diarrhoea.	Comment: Prevention of electrolyte depletion and replacement of electrolyte is 1st line treatment in acute diarrhoea. Oral rehydration therapy is listed in section 9.2. Codeine recommended only in short-term use due to CNS side effects and dependence.			
Antidiarrl	noeals			
	Codeine Phosphate	✓	✓	
	Loperamide (Note: - High doses may be used in Short Bowel Syndrome patients)	✓	*	

4. Exocrine	pancreatic insufficiency	Primary Care	Acute/ Specialist Services
Pancreatin			
	Creon [®]	\$	✓
	Pancrex V [®]	+	✓

5. Food Allergy - No formulary choices

6. Gastric acid disorders and ulceration	Primary Care	Acute/ Specialist Services	
Comment: Also refer to Forth Valley Dyspepsia Management Guidelines .			
Dyspepsia			
Antacids - Alginates			
Peptac (1 st line)	✓	✓	
(sodium alginate+calcium carbonate+sodium bicarbonate)			
Gaviscon Advance® (2 nd line)	✓	✓	
(sodium alginate + potassium bicarbonate)			
Infant Gaviscon® (magnesium alginate + sodium alginate)	✓	✓	
Aluminium and Magnesium containing antacids			
Co-magaldrox (Mucogel®)	✓	✓	

6.2 Gastri	c and Duodenal Ulceration	Primary Care	Acute/ Specialist Services
Gastropro	tective complexes and chelators		
	Sucralfate oral suspension 1g/5ml	+	✓
prescribed i	Sucralfate tablets are only available as an unlicensed special. License n preference to tablets. or antagonists	ed oral suspensi	ion should be
No formula			
Proton pu	mp inhibitors		
Comment:	Also refer to NHS Forth Valley - PPI Prescribing - Step Down in Lor	ng Term Use Gu	<u>iidance</u>
	Omeprazole Capsules (1 st line)	✓	✓
	Lansoprazole Capsules (2 nd line)	✓	✓
	Esomeprazole (3 rd line)	✓	✓
	Esomeprazole (I.v.)		✓
	Pantoprazole (i.v.)		✓

6.3 Gastro-oesophageal reflux disease	Primary Care	Acute/ Specialist Services	
Antacids - Alginates			
Peptac [®] (1 st line)	✓	✓	

Key \checkmark Initiate and continue \div Continue where appropriate.

(sodium alginate+calcium carbonate+sodium bicarbonate)		
--	--	--

7 Gastro-intestinal smooth muscle spasm		Primary Care	Acute/ Specialist Services
Antispasmodics			
	Mebeverine (not MR preparation) (1 st line)	✓	✓
	Hyoscine Butylbromide (2 nd line)	✓	✓

	er disorders and related conditions	Primary Care	Acute/ Specialist Services	
Bile a	Bile acids			
	Ursodeoxycholic acid (1 st line) (preferred formulation 250mg capsules)	\$	✓	
	Obeticholic acid (2 nd line)		✓	
Bile acid sequestrants				
	Colestyramine	+	✓	

9 Obesity		Primary Care	Acute/ Specialist Services
Li	iraglutide (Saxenda [®])	\$	✓
О	Orlistat	+	✓

Comment: Prescribing of Saxenda[®] and orlistat in Primary Care should only be undertaken following an initial assessment and recommendation by the FV Weight Management Service. Patients with BMI>30 with comorbidities as per SMC restrictions can be referred to the weight management service.

Liraglutide should be **prescribed as the brand name Saxenda**® if being prescribed for obesity management.

For Saxenda[®], in order to continue treatment patients must demonstrate at least a 5% weight loss from baseline after 3 months and a 10% weight loss after 12 months. Duration of treatment should be for a maximum of 2 years. Also see <u>SMC restrictions</u> for prescribing.

For further support patients should be signposted to the $\underline{\text{Choose to Lose Website}}$ and $\underline{\text{NHS Forth Valley weight}}$ $\underline{\text{management site}}$

10. Recta	l and anal disorders	Primary Care	Acute/ Specialist Services
Anal fissures			
	Glyceryl trinitrate rectal ointment 0.4% (Rectogesic®)	✓	✓
	Diltiazem 2% cream, ointment (unlicensed special)	 	✓

10.2 Pre	paration for Haemorrhoids	Primary Care	Acute/ Specialist Services
	Anusol [®]	✓	✓
	Anusol HC®	✓	✓
	Uniroid HC® (cinchocaine with hydrocortisone)	✓	✓
Comment: Uniroid HC [®] 1 st line if steroid containing preparation with anaesthetic is required, but steroid preparations are not 1 st line and should be used for no longer than seven days at a time.			
	Xyloproct® Ointment (hydrocortisone with lidocaine)	✓	✓
	Lidocaine (lignocaine) gel		✓

11.Stoma Care	Primary Care	Acute/ Specialist Services
Refer to NHS Forth Valley Stoma Formulary		